arXiv:1512.00136v1 [math.CO] 1 Dec 2015

A UNIVERSAL CLOSED FORM FOR SQUARE MATRIX POWERS

Walter Shur

20 Speyside Circle, Pittsboro, North Carolina, 27312, USA

Tel: 919-542-7179 Email: wrshur@gmail.com

Affiliation: None (Retired)

ABSTRACT.

This note presents a simple, universal closed form for the powers of any square matrix. A diligent search of the internet gave no indication that the form is known.

Keywords: Matrix, Powers, Closed Form

AMS Subject Classification: 15A99, 65F30

1. THE CLOSED FORM

Let $M = [m_{i,j}]$ be an n x n square matrix, real or complex, and let ${}_nm_{i,j}$ be the $(i,j)^{th}$ element of M^n. Then ${}_nm_{i,j}$ can be expressed as the sum of n terms, each of which is the product of a constant not depending on n, a binomial coefficient, and a power of an eigenvalue. In particular, for each unique eigenvalue, λ, of M, with multiplicity mp_λ, the portion of ${}_nm_{i,j}$ arising from λ is

$$c_{i,j,1}\tbinom{n-1}{0}\lambda^{n-1}+c_{i,j,2}\tbinom{n-1}{1}\lambda^{n-2}+\cdots+c_{i,j,mp_\lambda}\tbinom{n-1}{mp_\lambda-1}\lambda^{n-mp_\lambda}.$$

If the matrix is not singular, the formula works for any power of M, negative, zero or positive; if the matrix is singular, for positive powers only. In the case of zero eigenvalues, 0^n is taken as 0 if $n < 0$, and 1 if $n = 0$.

2. ILLUSTRATION

Let M be the matrix[1]

$$M = \begin{pmatrix} 66 & 83 & 95 & 31 & -50 & -63 \\ -71 & -79 & -86 & -22 & 59 & 72 \\ 83 & 74 & 69 & 4 & -77 & -90 \\ -74 & -50 & -34 & 16 & 77 & 86 \\ -7 & -31 & -47 & -32 & -12 & -5 \\ 65 & 89 & 105 & 41 & -40 & -56 \end{pmatrix}$$

The 6 eigenvalues of M are $\{-3, 2, 2, 1, 1, 1\}$. Hence,

$${}_nm_{i,j} = c_{i,j,1}(-3)^{n-1} + c_{i,j,2}2^{n-1} + c_{i,j,3}\tbinom{n-1}{1}2^{n-2} + c_{i,j,4}1^{n-1} + c_{i,j,5}\tbinom{n-1}{1}1^{n-2} + c_{i,j,6}\tbinom{n-1}{2}1^{n-3}.$$

To determine the values of the c constants for say, ${}_nm_{2,5}$, we first obtain by direct numerical calculation the values of ${}_nm_{2,5}$ for $n = 1$ to 6, namely, $59, 229, 764, 1915, 5270$, and 11377, respectively.

We then solve the 6 simultaneous equations

$${}_1m_{2,5} = 59,\ {}_2m_{2,5} = 229,\ {}_3m_{2,5} = 764,\ {}_4m_{2,5} = 1915,\ {}_5m_{2,5} = 5270,\ {}_6m_{2,5} = 11377,$$

obtaining

$$c_{2,5,1} = 3, c_{2,5,2} = 65, c_{2,5,3} = 126, c_{2,5,4} = -9, c_{2,5,5} = -9, c_{2,5,6} = 0.$$

Hence, we have the closed form

$${}_nm_{2,5} = 3(-3)^{n-1} + 65\ 2^{n-1} + 126\tbinom{n-1}{1}2^{n-2} - 9\ 1^{n-1} - 9\tbinom{n-1}{1}1^{n-2}.$$

3. DERIVATION OF THE CLOSED FORM

Any square matrix A can be written as $A = QTQ^{-1}$, where Q is a unitary matrix, and T is an upper triangular matrix (Schur decomposition). Hence, $A^n = QT^nQ^{-1}$. The eigenvalues of A and T are the same, and are the main diagonal elements of T. Theorem 3 in [2] shows that every element of T^n can be expressed by the linear combination described above in the first section of this note. Therefore, every

[1] this matrix appeared in [3]

element of A^n can also be expressed by that same general form.

4. CONCLUSION

The purpose of this note was not to present a new method of calculating the numerical values of the elements of a matrix raised to the n^{th} power. Nor does it present the only way to obtain a closed form for matrix powers. For example, the Mathematca[2] function MatrixPower[m,n][[2,5]] gives

$$ {}_nm_{2,5} = \tfrac{1}{2}(-2(-3)^n + 2^{1+n} - 18n + 63\ 2^n n). $$

This is identical to the closed form for ${}_nm_{2,5}$ given above, but with the beauty and symmetry absent.

Rather, the primary purpose of this note was to share with others the aesthetic beauty of a simple universal form, which represents some very complicated calculations.

References

[1] W. Shur, A Simple Closed Form For Triangular Matrix Powers. *Electronic Journal of Linear Algebra, 22:1000-1003,* 2011

[2] W. Shur, A Generalized Closed Form For Triangular Matrix Powers, *Cornell arXiv:1301.6820v2,* 2014

[3] H. E. Rose, Linear Algebra-A Pure Mathematical Approach, *ISBN 978-3-7643-6792-3,* 2002

[4] C.P. Huang, An Efficient Algorithm For Computing Powers Of Triangular Matrices, *ACM '78 Proceedings of the 1978 Annual Conference,* 2:954-957, 1978

11/30/15

[2] Version 10.3.0.0